Contents:

Volume 143, Number 1, October 5, 1998

In Brief

Multiple actin bundlers. Less order and more movement in the nucleus. Bax insertion into mitchondria. PtdIns(3,5)P_2 gets a kinase and a function.

W.A. Wells

Mini-Review

1 **Gene targeting studies begin to reveal the function of neurofilament proteins.**

N. Hirokawa and S. Takeda

Regular Articles

5 **Transcription sites are not correlated with chromosome territories in wheat nuclei.**

R. Abranches, A.F. Beven, L. Aragón-Alcaide, and P.J. Shaw

13 **Large-scale chromosomal movements during interphase progression in *Drosophila*.**

A.K. Csink and S. Henikoff

23 **Mutational analysis of the structure and localization of the nucleolus in the yeast *Saccharomyces cerevisiae*.**

M. Oakes, J.P. Aris, J.S. Brockenbrough, H. Wai, L. Vu, and M. Nomura

35 **The perinucleolar compartment and transcription.**

S. Huang, T.J. Deerinck, M.H. Ellisman, and D.L. Spector

49 **Characterization of the kinetochore binding domain of CENP-E reveals interactions with the kinetochore proteins CENP-F and hBUBR1.**

G.K.T. Chan, B.T. Schaar, and T.J. Yen

65 **Fab1p is essential for PtdIns(3)P 5-kinase activity and the maintenance of vacuolar size and membrane homeostasis.**

J.D. Gary, A.E. Wurmser, C.J. Bonangelino, L.S. Weisman, and S.D. Emr

81 **Sorting mechanisms regulating membrane protein traffic in the apical transcytotic pathway of polarized MDCK cells.**

A. Gibson, C.E. Futter, S. Maxwell, E.H. Allchin, M. Shipman, J.-P. Kraehenbuhl, D. Domingo, G. Odorizzi, I.S. Trowbridge, and C.R. Hopkins

95 **An atypical PKC directly associates and colocalizes at the epithelial tight junction with ASIP, a mammalian homologue of *Caenorhabditis elegans* polarity protein PAR-3.**

Y. Izumi, T. Hirose, Y. Tamai, S. Hirai, Y. Nagashima, T. Fujimoto, Y. Tabuse, K.J. Kemphues, and S. Ohno

107 **Small espin: A third actin-bundling protein and potential forked protein ortholog in brush border microvilli.**

J.R. Bartles, L. Zheng, A. Li, A. Wierda, and B. Chen

121 **Why are two different cross-linkers necessary for actin bundle formation in vivo and what does each cross-link contribute?**

L.G. Tilney, P.S. Connelly, K.A. Vranich, M.K. Shaw, and G.M. Guild

135 **Clb5-associated kinase activity is required early in the spindle pathway for correct preanaphase nuclear positioning in *Saccharomyces cerevisiae*.**

M. Segal, D.J. Clarke, and S.I. Reed

147 **Motile properties of vimentin intermediate filament networks in living cells.**

M. Yoon, R.D. Moir, V. Prahlad, and R.D. Goldman

159 **Rapid movements of vimentin on microtubule tracks: Kinesin-dependent assembly of intermediate filament networks.**

V. Prahlad, M. Yoon, R.D. Moir, R.D. Vale, and R.D. Goldman

171 **Neurofilament-dependent radial growth of motor axons and axonal organization of neurofilaments does not require the neurofilament heavy subunit (NF-H) or its phosphorylation.**

M.V. Rao, M.K. Houseweart, T.L. Williamson, T.O. Crawford, J. Folmer, and D.W. Cleveland

183 **Disruption of the NF-H gene increases axonal microtubule content and velocity of neurofilament transport: Relief of axonopathy resulting from the toxin β,β′-iminodipropionitrile.**

Q. Zhu, M. Lindenbaum, F. Levavasseur, H. Jacomy, and J.-P. Julien

195 **Requirement of heavy neurofilament subunit in the development of axons with large calibers.**

G.A. Elder, V.L. Friedrich, Jr., C. Kang, P. Bosco, A. Gourov, P.-H. Tu, B. Zhang, V.M.-Y. Lee, and R.A. Lazzarini

207 **Regulated targeting of BAX to mitochondria.**

I.S. Goping, A. Gross, J.N. Lavoie, M. Nguyen, R. Jemmerson, K. Roth, S.J. Korsmeyer, and G.C Shore

217 **Bax-induced cytochrome C release from mitochondria is independent of the permeability transition pore but highly dependent on Mg^{2+} ions.**

R. Eskes, B. Antonsson, A. Osen-Sand, S. Montessuit, C. Richter, R. Sadoul, G. Mazzei, A. Nichols, and J.-C. Martinou

225 **Transition from caspase-dependent to caspase-independent mechanisms at the onset of apoptotic execution.**
K. Samejima, S. Toné, T.J. Kottke, M. Enari, H. Sakahira, C.A. Cooke, F. Durrieu, L.M. Martins, S. Nagata, S.H. Kaufmann, and W.C. Earnshaw

241 **Modulation of calcium current in arteriolar smooth muscle by $\alpha_v\beta_3$ and $\alpha_5\beta_1$ integrin ligands.**
X. Wu, J.E. Mogford, S.H. Platts, G.E. Davis, G.A. Meininger, and M.J. Davis

253 **Cre-*loxP*–mediated inactivation of the α6A integrin splice variant in vivo: Evidence for a specific functional role of α6A in lymphocyte migration but not in heart development.**
C. Gimond, C. Baudoin, R. van der Neut, D. Kramer, J. Calafat, and A. Sonnenberg

267 **Fibronectin matrix regulates activation of RHO and CDC42 GTPases and cell cycle progression.**
S. Bourdoulous, G. Orend, D.A. MacKenna, R. Pasqualini, and E. Ruoslahti

277 ADDITIONS AND CORRECTIONS

Volume 143, Number 2, October 19, 1998

In Brief

Fusion and fission of mitochondria. How the extracellular matrix keeps cells alive. Another partner for talin. Releasing splicing factors.
W.A. Wells

Commentary

279 **Growth factors in the nucleolus?**
T. Pederson

Regular Articles

283 **Spindle checkpoint protein Xmad1 recruits Xmad2 to unattached kinetochores.**
R.-H. Chen, A. Shevchenko, M. Mann, and A.W. Murray

297 **Serine phosphorylation of SR proteins is required for their recruitment to sites of transcription in vivo.**
T. Misteli, J.F. Cáceres, J.Q. Clement, A.R. Krainer, M.F. Wilkinson, and D.L. Spector

309 **Determination of the functional domain organization of the importin α nuclear import factor.**
A. Herold, R. Truant, H. Wiegand, and B.R. Cullen

319 **The membrane transport factor TAP/p115 cycles between the Golgi and earlier secretory compartments and contains distinct domains required for its localization and function.**
D.S. Nelson, C. Alvarez, Y.-s. Gao, R. García-Mata, E. Fialkowski, and E. Sztul

333 **The dynamin-related GTPase, Dnm1p, controls mitochondrial morphology in yeast.**
D. Otsuga, B.R. Keegan, E. Brisch, J.W. Thatcher, G.J. Hermann, W. Bleazard, and J.M. Shaw

351 **A human dynamin-related protein controls the distribution of mitochondria.**
E. Smirnova, D.-L. Shurland, S.N. Ryazantsev, and A.M. van der Bliek

359 **Mitochondrial fusion in yeast requires the transmembrane GTPase Fzo1p.**
G.J. Hermann, J.W. Thatcher, J.P. Mills, K.G. Hales, M.T. Fuller, J. Nunnari, and J.M. Shaw

375 **Identification of Kel1p, a Kelch domain-containing protein involved in cell fusion and morphology in *Saccharomyces cerevisiae*.**
J. Philips and I. Herskowitz

391 **A single gene product, claudin-1 or -2, reconstitutes tight junction strands and recruits occludin in fibroblasts.**
M. Furuse, H. Sasaki, K. Fujimoto, and S. Tsukita

403 **CDO, a Robo-related cell surface protein that mediates myogenic differentiation.**
J.-S. Kang, P.J. Mulieri, C. Miller, D.A. Sassoon, and R.S. Krauss

415 **imp2, a new component of the actin ring in the fission yeast *Schizosaccharomyces pombe*.**
J. Demeter and S. Sazer

429 **Layilin, a novel talin-binding transmembrane protein homologous with C-type lectins, is localized in membrane ruffles.**
M.L. Borowsky and R.O. Hynes

443 **Suppression of radixin and moesin alters growth cone morphology, motility, and process formation in primary cultured neurons.**
G. Paglini, P. Kunda, S. Quiroga, K. Kosik, and A. Cáceres

457 **Cyclin-dependent kinases participate in death of neurons evoked by DNA-damaging agents.**
D.S. Park, E.J. Morris, J. Padmanabhan, M.L. Shelanski, H.M. Geller, and L.A. Greene

469 **Onset of keratin 17 expression coincides with the definition of major epithelial lineages during skin development.**
K.M. McGowan and P.A. Coulombe

487 **Functional differences between keratins of stratified and simple epithelia.**
E. Hutton, R.D. Paladini, Q.-C. Yu, M. Yen, P.A. Coulombe, and E. Fuchs

501 **Visualization of phosphoinositides that bind pleckstrin homology domains: Calcium- and agonist-induced dynamic changes and relationship to Myo-[^{3}H]inositol-labeled phosphoinositide pools.**
P. Várnai and T. Balla

511 **Distinct effects of fatty acids on translocation of γ- and ϵ-subspecies of protein kinase C.**
Y. Shirai, K. Kashiwagi, K. Yagi, N. Sakai, and N. Saito

523 **The nonreceptor protein tyrosine phosphatase PTP1B binds to the cytoplasmic domain of N-cadherin and regulates the cadherin–actin linkage.**
J. Balsamo, C. Arregui, T. Leung, and J. Lilien

533 **Reconstitution of mammary gland development in vitro: Requirement of c-met and c-erbB2 signaling for branching and alveolar morphogenesis.**
C. Niemann, V. Brinkmann, E. Spitzer, G. Hartmann, M. Sachs, H. Naundorf, and W. Birchmeier

547 **Extracellular matrix survival signals transduced by focal adhesion kinase suppress p53-mediated apoptosis.**
D. Ilić, E.A.C. Almeida, D.D. Schlaepfer, P. Dazin, S. Aizawa, and C.H. Damsky

561 ADDITIONS AND CORRECTIONS

Volume 143, Number 3, November 2, 1998

In Brief

Rethinking septin function. Making space in the extracellular matrix. CFTR's chloride sensor, not conductance, triggers ATP release. Immunosuppression by disrupting rafts.
W.A. Wells

Regular Articles

563 **Recruitment of phosphorylated chromatin assembly factor 1 to chromatin after UV irradiation of human cells.**
E. Martini, D.M.J. Roche, K. Marheineke, A. Verreault, and G. Almouzni

577 **Molecular architecture of the yeast nuclear pore complex: Localization of Nsp1p subcomplexes.**
B. Fahrenkrog, E.C. Hurt, U. Aebi, and N. Panté

589 **Reconstitution of retrograde transport from the Golgi to the ER in vitro.**
A. Spang and R. Schekman

601 **KDEL receptor (Erd2p)-mediated retrograde transport of the cholera toxin A subunit from the Golgi involves COPI, p23, and the COOH terminus of Erd2p.**
I. Majoul, K. Sohn, F.T. Wieland, R. Pepperkok, M. Pizza, J. Hillemann, and H.-D. Söling

613 **ZAP-70 association with T cell receptor ζ (TCRζ): Fluorescence imaging of dynamic changes upon cellular stimulation.**
J. Sloan-Lancaster, J. Presley, J. Ellenberg, T. Yamazaki, J. Lippincott-Schwartz, and L.E. Samelson

625 **The UDP-Glc:glycoprotein glucosyltransferase is essential for *Schizosaccharomyces pombe* viability under conditions of extreme endoplasmic reticulum stress.**
S. Fanchiotti, F. Fernández, C. D'Alessio, and A.J. Parodi

637 **Polyunsaturated fatty acids inhibit T cell signal transduction by modification of detergent-insoluble membrane domains.**
T.M. Stulnig, M. Berger, T. Sigmund, D. Raederstorff, H. Stockinger, and W. Waldhäusl

645 **Cystic fibrosis transmembrane conductance regulator–associated ATP release is controlled by a chloride sensor.**
Q. Jiang, D. Mak, S. Devidas, E.M. Schwiebert, A. Bragin, Y. Zhang, W.R. Skach, W.B. Guggino, J.K. Foskett, and J.F. Engelhardt

659 ***Drosophila* polo kinase is required for cytokinesis.**
M. Carmena, M.G. Riparbelli, G. Minestrini, Á.M. Tavares, R. Adams, G. Callaini, and D.M. Glover

673 **Localization of the kinesin-like protein Xklp2 to spindle poles requires a leucine zipper, a microtubule-associated protein, and dynein.**
T. Wittmann, H. Boleti, C. Antony, E. Karsenti, and I. Vernos

687 **Time-lapse microscopy reveals unique roles for kinesins during anaphase in budding yeast.**
A.F. Straight, J.W. Sedat, and A.W. Murray

695 **Novel features of the light chain of microtubule-associated protein MAP1B: Microtubule stabilization, self interaction, actin filament binding, and regulation by the heavy chain.**
M. Tögel, G. Wiche, and F. Propst

709 **The septins are required for the mitosis-specific activation of the Gin4 kinase.**
C.W. Carroll, R. Altman, D. Schieltz, J.R. Yates III, and D. Kellogg

719 **Role of the yeast Gin4p protein kinase in septin assembly and the relationship between septin assembly and septin function.**
M.S. Longtine, H. Fares, and J.R. Pringle

737 **Polymerization of purified yeast septins: Evidence that organized filament arrays may not be required for septin function.**
J.A. Frazier, M.L. Wong, M.S. Longtine, J.R. Pringle, M. Mann, T.J. Mitchison, and C. Field

751 **The yeast centrin, Cdc31p, and the interacting protein kinase, Kic1p, are required for cell integrity.**
D.S. Sullivan, S. Biggins, and M.D. Rose

767 **Cell cycle–related changes in the conducting properties of r-eag K^+ channels.**
L.A. Pardo, A. Brüggemann, J. Camacho, and W. Stühmer

777 **Overexpression of tau protein inhibits kinesin-dependent trafficking of vesicles, mitochondria, and endoplasmic reticulum: Implications for Alzheimer's disease.**
A. Ebneth, R. Godemann, K. Stamer, S. Illenberger, B. Trinczek, E.-M. Mandelkow, and E. Mandelkow

795 **Paralemmin, a prenyl-palmitoyl–anchored phosphoprotein abundant in neurons and implicated in plasma membrane dynamics and cell process formation.**
C. Kutzleb, G. Sanders, R. Yamamoto, X. Wang, B. Lichte, E. Petrasch-Parwez, and M.W. Kilimann

815 **Targeted ablation of connexin50 in mice results in microphthalmia and zonular pulverulent cataracts.**
T.W. White, D.A. Goodenough, and D.L. Paul

827 **TGF-β plays a key role in morphogenesis of the pancreatic islets of Langerhans by controlling the activity of the matrix metalloproteinase MMP-2.**
F. Miralles, T. Battelino, P. Czernichow, and R. Scharfmann

837 **CD44 occupancy prevents macrophage multinucleation.**
H. Sterling, C. Saginario, and A. Vignery

849 **Dystrophic muscle in mice chimeric for expression of α5 integrin.**
D. Taverna, M.-H. Disatnik, H. Rayburn, R.T. Bronson, J. Yang, T.A. Rando, and R.O. Hynes

861 **Impaired integrin-mediated adhesion and signaling in fibroblasts expressing a dominant-negative mutant PTP1B.**
C.O. Arregui, J. Balsamo, and J. Lilien

Volume 143, Number 4, November 16, 1998

In Brief

A profusion of sorting pathways. Nuclear import without importins. Kinesin inhibition.
W.A. Wells

Regular Articles

875 **Characterization of HIV-1 Vpr nuclear import: Analysis of signals and pathways.**
Y. Jenkins, M. McEntee, K. Weis, and W.C. Greene

887 **Nuclear import and the evolution of a multifunctional RNA-binding protein.**
J.S. Rosenblum, L.F. Pemberton, N. Bonifaci, and G. Blobel

901 **In vivo function of Hsp90 is dependent on ATP binding and ATP hydrolysis.**
W.M.J. Obermann, H. Sondermann, A.A. Russo, N.P. Pavletich, and F.U. Hartl

911 **Changes in organization of *Crithidia fasciculata* kinetoplast DNA replication proteins during the cell cycle.**
C.E. Johnson and P.T. Englund

921 **A role for the DnaJ homologue Scj1p in protein folding in the yeast endoplasmic reticulum.**
S. Silberstein, G. Schlenstedt, P.A. Silver, and R. Gilmore

935 **The high osmolarity glycerol response (HOG) MAP kinase pathway controls localization of a yeast golgi glycosyltransferase.**
T.B. Reynolds, B.D. Hopkins, M.R. Lyons, and T.R. Graham

947 **Neuroendocrine synaptic vesicles are formed in vitro by both clathrin-dependent and clathrin-independent pathways.**
G. Shi, V. Faúndez, J. Roos, E.C. Dell'Angelica, and R.B. Kelly

957 **Syntaxin 13 mediates cycling of plasma membrane proteins via tubulovesicular recycling endosomes.**
R. Prekeris, J. Klumperman, Y.A. Chen, and R.H. Scheller

973 **Direct pathway from early/recycling endosomes to the golgi apparatus revealed through the study of Shiga toxin B-fragment transport.**
F. Mallard, C. Antony, D. Tenza, J. Salamero, B. Goud, and L. Johannes

991 **Tic20 and Tic22 are new components of the protein import apparatus at the chloroplast inner envelope membrane.**
A. Kouranov, X. Chen, B. Fuks, and D.J. Schnell

1003 **Interferon α inhibits a Src-mediated pathway necessary for *Shigella*-induced cytoskeletal rearrangements in epithelial cells.**
G. Duménil, J.C. Olivo, S. Pellegrini, M. Fellous, P.J. Sansonetti, and G. Tran Van Nhieu

1013 **The NH_2 terminus of titin spans the Z-disc: Its interaction with a novel 19-kD ligand (T-cap) is required for sarcomeric integrity.**
C.C. Gregorio, K. Trombitás, T. Centner, B. Kolmerer, G. Stier, K. Kunke, K. Suzuki, F. Obermayr, B. Herrmann, H. Granzier, H. Sorimachi, and S. Labeit

1029 ***Saccharomyces cerevisiae* Duo1p and Dam1p, novel proteins involved in mitotic spindle function.**
C. Hofmann, I.M. Cheeseman, B.L. Goode, K.L. McDonald, G. Barnes, and D.G. Drubin

1041 **When overexpressed, a novel centrosomal protein, RanBPM, causes ectopic microtubule nucleation similar to γ-tubulin.**
M. Nakamura, H. Masuda, J. Horii, K.-i. Kuma, N. Yokoyama, T. Ohba, H. Nishitani, T. Miyata, M. Tanaka, and T. Nishimoto

1053 **Light chain–dependent regulation of kinesin's interaction with microtubules.**
K.J. Verhey, D.L. Lizotte, T. Abramson, L. Barenboim, B.J. Schnapp, and T.A. Rapoport

1067 **A functional role for specific spliced variants of the α7β1 integrin in acetylcholine receptor clustering.**
D.J. Burkin, M. Gu, B.L. Hodges, J.T. Campanelli, and S.J. Kaufman

1077 **Dynamics of axonal microtubules regulate the topology of new membrane insertion into the growing neurites.**
S. Zakharenko and S. Popov

1087 **The transcription factor AP-1 is required for EGF-induced activation of Rho-like GTPases, cytoskeletal rearrangements, motility, and in vitro invasion of A431 cells.**
A. Malliri, M. Symons, R.F. Hennigan, A.F.L. Hurlstone, R.F. Lamb, T. Wheeler, and B.W. Ozanne

1101 **Identification of a bipotential precursor cell in hepatic cell lines derived from transgenic mice expressing cyto-met in the liver.**
F.M. Spagnoli, L. Amicone, M. Tripodi, and M.C. Weiss

1113 **Osteoblast recruitment and bone formation enhanced by cell matrix–associated heparin-binding growth-associated molecule (HB-GAM).**
S. Imai, M. Kaksonen, E. Raulo, T. Kinnunen, C. Fages, X. Meng, M. Lakso, and H. Rauvala

1129 **Role of P-selectin cytoplasmic domain in granular targeting in vivo and in early inflammatory responses.**
D.W. Hartwell, T.N. Mayadas, G. Berger, P.S. Frenette, H. Rayburn, R.O. Hynes, and D.D. Wagner

1143 ADDITIONS AND CORRECTIONS

Volume 143, Number 5, November 30, 1998

In Brief

Kakapo keeps it all together. Collagen squeezes out the notochord. Viral fusion machinery. Multiple targets for PTEN. For Tiam1, location is everything.
W.A. Wells

Regular Articles

1145 **Processing of the *psbA* 5′ untranslated region in *Chlamydomonas reinhardtii* depends upon factors mediating ribosome association.**
R.K. Bruick and S.P. Mayfield

1155 **Membrane fusion mediated by baculovirus gp64 involves assembly of stable gp64 trimers into multiprotein aggregates.**
I. Markovic, H. Pulyaeva, A. Sokoloff, and L.V. Chernomordik

1167 **Involvement of long chain fatty acid elongation in the trafficking of secretory vesicles in yeast.**
D. David, S. Sundararababu, and J.E. Gerst

1183 **Integral membrane protein sorting to vacuoles in plant cells: Evidence for two pathways.**
L. Jiang and J.C. Rogers

1201 **Ca^{2+}-dependent muscle dysfunction caused by mutation of the *Caenorhabditis elegans* troponin T-1 gene.**
K. McArdle, T.StC. Allen, and E.A. Bucher

1215 ***Unc-45* mutations in *Caenorhabditis elegans* implicate a CRO1/She4p-like domain in myosin assembly.**
J.M. Barral, C.C. Bauer, I. Ortiz, and H.F. Epstein

1227 **Nexilin: A novel actin filament-binding protein localized at cell–matrix adherens junction.**
T. Ohtsuka, H. Nakanishi, W. Ikeda, A. Satoh, Y. Momose, H. Nishioka, and Y. Takai

1239 **The "8-kD" cytoplasmic dynein light chain is required for nuclear migration and for dynein heavy chain localization in *Aspergillus nidulans*.**
S.M. Beckwith, C.H. Roghi, B. Liu, and N.R. Morris

1249 **Roles of Rho-associated kinase in cytokinesis; mutations in Rho-associated kinase phosphorylation sites impair cytokinetic segregation of glial filaments.**
Y. Yasui, M. Amano, K.-i. Nagata, N. Inagaki, H. Nakamura, H. Saya, K. Kaibuchi, and M. Inagaki

1259 **Kakapo, a novel cytoskeletal-associated protein is essential for the restricted localization of the neuregulin-like factor, Vein, at the muscle–tendon junction site.**
D. Strumpf and T. Volk

1271 ***kakapo*, a gene required for adhesion between and within cell layers in *Drosophila*, encodes a large cytoskeletal linker protein related to plectin and dystrophin.**
S.L. Gregory and N.H. Brown

1283 **The *kakapo* mutation affects terminal arborization and central dendritic sprouting of *Drosophila* motorneurons.**
A. Prokop, J. Uhler, J. Roote, and M. Bate

1295 **Ankyrin$_G$ is required for clustering of voltage-gated Na channels at axon initial segments and for normal action potential firing.**
D. Zhou, S. Lambert, P.L. Malen, S. Carpenter, L.M. Boland, and V. Bennett

1305 **Nervous system defects of ankyrin$_B$ (−/−) mice suggest functional overlap between the cell adhesion molecule L1 and 440-kD ankyrin$_B$ in premyelinated axons.**
P. Scotland, D. Zhou, H. Benveniste, and V. Bennett

1317 **Removal of the membrane-anchoring domain of epidermal growth factor leads to intracrine signaling and disruption of mammary epithelial cell organization.**
H.S. Wiley, M.F. Woolf, L.K. Opresko, P.M. Burke, B. Will, J.R. Morgan, and D.A. Lauffenburger

1329 **In vitro repression of Brca1-associated RING domain gene, *Bard1*, induces phenotypic changes in mammary epithelial cells.**
I. Irminger-Finger, J.V. Soriano, G. Vaudan, R. Montesano, and A.-P. Sappino

1341 **Integration of endothelial cells in multicellular spheroids prevents apoptosis and induces differentiation.**
T. Korff and H.G. Augustin

1353 **Caspase-independent cell killing by Fas-associated protein with death domain.**
A. Kawahara, Y. Ohsawa, H. Matsumura, Y. Uchiyama, and S. Nagata

1361 **SAPK2/p38-dependent F-actin reorganization regulates early membrane blebbing during stress-induced apoptosis.**
J. Huot, F. Houle, S. Rousseau, R.G. Deschesnes, G.M. Shah, and J. Landry

1375 **Tumor suppressor PTEN inhibits integrin- and growth factor–mediated mitogen-activated protein (MAP) kinase signaling pathways.**
J. Gu, M. Tamura, and K.M. Yamada

1385 **Matrix-dependent Tiam1/Rac signaling in epithelial cells promotes either cell–cell adhesion or cell migration and is regulated by phosphatidylinositol 3-kinase.**
E.E. Sander, S. van Delft, J.P. ten Klooster, T. Reid, R.A. van der Kammen, F. Michiels, and J.G. Collard

1399 **Collagen II is essential for the removal of the notochord and the formation of intervertebral discs.**
A. Aszódi, D. Chan, E. Hunziker, J.F. Bateman, and R. Fässler

1413 ADDITIONS AND CORRECTIONS

Volume 143, Number 6, December 14, 1998

In Brief

Membrane trafficking by Kinesin II. Axonin-1 as an accessory adhesion protein. Paired pathways.
W.A. Wells

Regular Articles

1415 **Spatial and temporal dynamics of DNA replication sites in mammalian cells.**
H. Ma, J. Samarabandu, R.S. Devdhar, R. Acharya, P.-c. Cheng, C. Meng, and R. Berezney

1427 **Chromosomal proteins HMG-14 and HMG-17 are released from mitotic chromosomes and imported into the nucleus by active transport.**
R. Hock, U. Scheer, and M. Bustin

1437 **pEg7, a new *Xenopus* protein required for mitotic chromosome condensation in egg extracts.**
F. Cubizolles, V. Legagneux, R. Le Guellec, I. Chartrain, R. Uzbekov, C. Ford, and K. Le Guellec

1447 **A novel nuclear import pathway for the transcription factor TFIIS.**
M. Albertini, L.F. Pemberton, J.S. Rosenblum, and G. Blobel

1457 **Recruitment and the role of nuclear localization in polyglutamine-mediated aggregation.**
M.K. Perez, H.L. Paulson, S.J. Pendse, S.J. Saionz, N.M. Bonini, and R.N. Pittman

1471 **Interaction of ZPR1 with translation elongation factor-1α in proliferating cells.**
L. Gangwani, M. Mikrut, Z. Galcheva-Gargova, and R.J. Davis

1485 **Kinetic analysis of secretory protein traffic and characterization of Golgi to plasma membrane transport intermediates in living cells.**
K. Hirschberg, C.M. Miller, J. Ellenberg, J.F. Presley, E.D. Siggia, R.D. Phair, and J. Lippincott-Schwartz

1505 **Recycling of Golgi-resident glycosyltransferases through the ER reveals a novel pathway and provides an explanation for nocodazole-induced Golgi scattering.**
B. Storrie, J. White, S. Röttger, E.H.K. Stelzer, T. Suganuma, and T. Nilsson

1523 **Two functional states of the CD11b A-domain: Correlations with key features of two Mn^{2+}-complexed crystal structures.**
R. Li, P. Rieu, D.L. Griffith, D. Scott, and M.A. Arnaout

1535 **The localization of myosin VI at the Golgi complex and leading edge of fibroblasts and its phosphorylation and recruitment into membrane ruffles of A431 cells after growth factor stimulation.**
F. Buss, J. Kendrick-Jones, C. Lionne, A.E. Knight, G.P. Côté, and J.P. Luzio

1547 **Heterotrimeric kinesin II is the microtubule motor protein responsible for pigment dispersion in *Xenopus* melanophores.**
M.C. Tuma, A. Zill, N. Le Bot, I. Vernos, and V. Gelfand

1559 **Role of Xklp3, a subunit of the *Xenopus* Kinesin II heterotrimeric complex, in membrane transport between the endoplasmic reticulum and the Golgi apparatus.**
N. Le Bot, C. Antony, J. White, E. Karsenti, and I. Vernos

1575 **Centriole disassembly in vivo and its effect on centrosome structure and function in vertebrate cells.**
Y. Bobinnec, A. Khodjakov, L.M. Mir, C.L. Rieder, B. Eddé, and M. Bornens

1591 **Distinct mutants of retrograde intraflagellar transport (IFT) share similar morphological and molecular defects.**
G. Piperno, E. Siuda, S. Henderson, M. Segil, H. Vaananen, and M. Sassaroli

1603 **Role of polo kinase and Mid1p in determining the site of cell division in fission yeast.**
J. Bähler, A.B. Steever, S. Wheatley, Y.-l. Wang, J.R. Pringle, K.L. Gould, and D. McCollum

1617 **Morphogenesis beyond cytokinetic arrest in *Saccharomyces cerevisiae*.**
J. Jiménez, V.J. Cid, R. Cenamor, M. Yuste, G. Molero, C. Nombela, and M. Sánchez

1635 **AIR-2: An Aurora/Ipl1-related protein kinase associated with chromosomes and midbody microtubules is required for polar body extrusion and cytokinesis in *Caenorhabditis elegans* embryos.**
J.M. Schumacher, A. Golden, and P.J. Donovan

1647 **Distinct roles for the p110α and hVPS34 phosphatidylinositol 3′-kinases in vesicular trafficking, regulation of the actin cytoskeleton, and mitogenesis.**
U. Siddhanta, J. McIlroy, A. Shah, Y. Zhang, and J.M. Backer

1661 **Calcium and protein kinase C regulate the actin cytoskeleton in the synaptic terminal of retinal bipolar cells.**
C. Job and L. Lagnado

1673 **Neurite fasciculation mediated by complexes of axonin-1 and Ng cell adhesion molecule.**
S. Kunz, M. Spirig, C. Ginsburg, A. Buchstaller, P. Berger, R. Lanz, C. Rader, L. Vogt, B. Kunz, and P. Sonderegger

1691 **P53 is essential for developmental neuron death as regulated by the TrkA and p75 neurotrophin receptors.**
R.S. Aloyz, S.X. Bamji, C.D. Pozniak, J.G. Toma, J. Atwal, D.R. Kaplan, and F.D. Miller

1705 **Overexpression of VEGF in testis and epididymis causes infertility in transgenic mice: Evidence for nonendothelial targets for VEGF.**
E.I. Korpelainen, M.J. Karkkainen, A. Tenhunen, M. Lakso, H. Rauvala, M. Vierula, M. Parvinen, and K. Alitalo

1713 **Roles for laminin in embryogenesis: Exencephaly, syndactyly, and placentopathy in mice lacking the laminin α5 chain.**
J.H. Miner, J. Cunningham, and J.R. Sanes

1725 **Gap junction–mediated cell–cell communication modulates mouse neural crest migration.**
G.Y. Huang, E.S. Cooper, K. Waldo, M.L. Kirby, N.B. Gilula, and C.W. Lo

1735 **Cellular interaction of integrin α3β1 with laminin 5 promotes gap junctional communication.**
P.D. Lampe, B.P. Nguyen, S. Gil, M. Usui, J. Olerud, Y. Takada, and W.G. Carter

1749 **Release of cAMP gating by the α6β4 integrin stimulates lamellae formation and the chemotactic migration of invasive carcinoma cells.**
K.L. O'Connor, L.M. Shaw, and A.M. Mercurio

1761 ADDITIONS AND CORRECTIONS

In Brief

Movement by cable or capture. Gathering together unfolded proteins. Not all sarcoglycans are equal. Nuclear pore complexes and spindle pole bodies share a component. Early functions for desmosomes. A sex-specific homeodomain protein in algae.

W.A. Wells

Regular Articles

1763 **INCENP centromere and spindle targeting: Identification of essential conserved motifs and involvement of heterochromatin protein HP1.**

A.M. Ainsztein, S.E. Kandels-Lewis, A.M. Mackay, and W.C. Earnshaw

1775 **Fission yeast Bub1 is a mitotic centromere protein essential for the spindle checkpoint and the preservation of correct ploidy through mitosis.**

P. Bernard, K. Hardwick, and J.-P. Javerzat

1789 ***Saccharomyces cerevisiae* Ndc1p is a shared component of nuclear pore complexes and spindle pole bodies.**

H.J. Chial, M.P. Rout, T.H. Giddings, Jr., and M. Winey

1801 **Functional analysis of Tpr: Identification of nuclear pore complex association and nuclear localization domains and a role in mRNA export.**

P. Bangs, B. Burke, C. Powers, R. Craig, A. Purohit, and S. Doxsey

1813 **Specific binding of the karyopherin Kap121p to a subunit of the nuclear pore complex containing Nup53p, Nup59p, and Nup170p.**

M. Marelli, J.D. Aitchison, and R.W. Wozniak

1831 **Homotypic fusion of immature secretory granules during maturation in a cell-free assay.**

S. Urbé, L.J. Page, and S.A. Tooze

1845 **Biochemical and functional studies of cortical vesicle fusion: The SNARE complex and Ca^{2+} sensitivity.**

J.R. Coorssen, P.S. Blank, M. Tahara, and J. Zimmerberg

1859 **Pex18p and Pex21p, a novel pair of related peroxins essential for peroxisomal targeting by the PTS2 pathway.**

P.E. Purdue, X. Yang, and P.B. Lazarow

1871 **Redundant and distinct functions for dynamin-1 and dynamin-2 isoforms.**

Y. Altschuler, S.M. Barbas, L.J. Terlecky, K. Tang, S. Hardy, K.E. Mostov, and S.L. Schmid

1883 **Aggresomes: A cellular response to misfolded proteins.**

J.A. Johnston, C.L. Ward, and R.R. Kopito

1899 **Visualization of melanosome dynamics within wild-type and dilute melanocytes suggests a paradigm for myosin V function in vivo.**

X. Wu, B. Bowers, K. Rao, Q. Wei, and J.A. Hammer III

1919 **Visualization and molecular analysis of actin assembly in living cells.**

D.A. Schafer, M.D. Welch, L.M. Machesky, P.C. Bridgman, S.M. Meyer, and J.A. Cooper

1931 **Tropomyosin-containing actin cables direct the Myo2p-dependent polarized delivery of secretory vesicles in budding yeast.**

D.W. Pruyne, D.H. Schott, and A. Bretscher

1947 **Dual function of Cyk2, a cdc15/PSTPIP family protein, in regulating actomyosin ring dynamics and septin distribution.**

J. Lippincott and R. Li

1961 **Nonuniform microtubular polarity established by CHO1/MKLP1 motor protein is necessary for process formation of podocytes.**

N. Kobayashi, J. Reiser, W. Kriz, R. Kuriyama, and P. Mundel

1971 **A gamete-specific, sex-limited homeodomain protein in *Chlamydomonas*.**

V. Kurvari, N.V. Grishin, and W.J. Snell

1981 **ARF1 mediates paxillin recruitment to focal adhesions and potentiates Rho-stimulated stress fiber formation in intact and permeabilized Swiss 3T3 fibroblasts.**

J.C. Norman, D. Jones, S.T. Barry, M.R. Holt, S. Cockcroft, and D.R. Critchley

1997 **Regulation of the cell cycle by focal adhesion kinase.**

J.-H. Zhao, H. Reiske, and J.-L. Guan

2009 **Desmoplakin is required early in development for assembly of desmosomes and cytoskeletal linkage.**

G.I. Gallicano, P. Kouklis, C. Bauer, M. Yin, V. Vasioukhin, L. Degenstein, and E. Fuchs

2023 **Mutation of a major keratin phosphorylation site predisposes to hepatotoxic injury in transgenic mice.**

N.-O. Ku, S.A. Michie, R.M. Soetikno, E.Z. Resurreccion, R.L. Broome, and M.B. Omary

2033 **Molecular organization of sarcoglycan complex in mouse myotubes in culture.**

Y.-m. Chan, C.G. Bönnemann, H.G.W. Lidov, and L.M. Kunkel

2045 **Functional characteristics of ES cell–derived cardiac precursor cells identified by tissue-specific expression of the green fluorescent protein.**

E. Kolossov, B.K. Fleischmann, Q. Liu, W. Bloch, S. Viatchenko-Karpinski, O. Manzke, G.J. Ji, H. Bohlen, K. Addicks, and J. Hescheler

2057 **Autocrine tumor necrosis factor (TNF) and lymphotoxin (LT) α differentially modulate cellular sensitivity to TNF/LT-α cytotoxicity in L929 cells.**
E. Decoster, S. Cornelis, B. Vanhaesebroeck, and W. Fiers

2067 **The cell adhesion molecule L1 is developmentally regulated in the renal epithelium and is involved in kidney branching morphogenesis.**
H. Debiec, E.I. Christensen, and P.M. Ronco

2081 **Activation of $\alpha_V\beta_3$ on vascular cells controls recognition of prothrombin.**
T.V. Byzova and E.F. Plow

2093 ADDITIONS AND CORRECTIONS

2095 AUTHOR INDEX FOR VOLUMES 140–143

2131 SUBJECT INDEX FOR VOLUMES 140–143

2157 ACKNOWLEDGMENT